


# PALS Systematic Approach Algorithm



American Academy of Pediatrics  
DEDICATED TO THE HEALTH OF ALL CHILDREN™

## Pediatric Advanced Life Support



# Pediatric Septic Shock Algorithm




American Academy  
of Pediatrics



DEDICATED TO THE HEALTH OF ALL CHILDREN™

## Pediatric Advanced Life Support



Modified from Brierley J, Carcillo JA, Choong K, Cornell T, Decaen A, Deymann A, Doctor A, Davis A, Duff J, Dugas MA, Duncan A, Evans B, Feldman J, Felmet K, Fisher G, Frankel L, Jeffries H, Greenwald B, Gutierrez J, Hall M, Han YY, Hanson J, Hazelzet J, Hernan L, Kiff J, Kissoon N, Kon A, Irazusta J, Lin J, Lorts A, Mariscalco M, Mehta R, Nadel S, Nguyen T, Nicholson C, Peters M, Okhuysen-Cawley R, Poulton T, Relves M, Rodriguez A, Rozenfeld R, Schnitzler E, Shanley T, Kache S, Skippen P, Torres A, von Dessauer B, Weingarten J, Yeh T, Zaritsky A, Stojadinovic B, Zimmerman J, Zuckerberg A. Clinical practice parameters for hemodynamic support of pediatric and neonatal septic shock: 2007 update from the American College of Critical Care Medicine. *Crit Care Med*. 2009;37(2):666-688.





# Pediatric Bradycardia With a Pulse and Poor Perfusion Algorithm




American Heart Association®

American Academy of Pediatrics



DEDICATED TO THE HEALTH OF ALL CHILDREN™

## Pediatric Advanced Life Support



# Pediatric Tachycardia With a Pulse and Adequate Perfusion Algorithm




American Academy  
of Pediatrics



DEDICATED TO THE HEALTH OF ALL CHILDREN™

## Pediatric Advanced Life Support





# Pediatric Tachycardia With a Pulse and Poor Perfusion Algorithm




American Academy of Pediatrics



DEDICATED TO THE HEALTH OF ALL CHILDREN™

## Pediatric Advanced Life Support



# Pediatric Tachycardia With a Pulse and Poor Perfusion Algorithm




American Academy of Pediatrics



DEDICATED TO THE HEALTH OF ALL CHILDREN™

## Pediatric Advanced Life Support





# Pediatric Cardiac Arrest Algorithm




American Academy  
of Pediatrics



DEDICATED TO THE HEALTH OF ALL CHILDREN™

## Pediatric Advanced Life Support

### Shout for Help/Activate Emergency Response



### Doses/Details

#### CPR Quality

- Push hard ( $\geq 1/3$  of anterior-posterior diameter of chest) and fast (at least 100/min) and allow complete chest recoil
- Minimize interruptions in compressions
- Avoid excessive ventilation
- Rotate compressor every 2 minutes
- If no advanced airway, 15:2 compression-ventilation ratio. If advanced airway, 8-10 breaths per minute with continuous chest compressions

#### Shock Energy for Defibrillation

First shock 2 J/kg, second shock 4 J/kg, subsequent shocks  $\geq 4$  J/kg, maximum 10 J/kg or adult dose.

#### Drug Therapy

- **Epinephrine IO/IV Dose:** 0.01 mg/kg (0.1 mL/kg of 1:10 000 concentration). Repeat every 3-5 minutes. If no IO/IV access, may give endotracheal dose: 0.1 mg/kg (0.1 mL/kg of 1:1000 concentration).
- **Amiodarone IO/IV Dose:** 5 mg/kg bolus during cardiac arrest. May repeat up to 2 times for refractory VF/pulseless VT.

#### Advanced Airway

- Endotracheal intubation or supraglottic advanced airway
- Waveform capnography or capnometry to confirm and monitor ET tube placement
- Once advanced airway in place give 1 breath every 6-8 seconds (8-10 breaths per minute)

#### Return of Spontaneous Circulation (ROSC)

- Pulse and blood pressure
- Spontaneous arterial pressure waves with intra-arterial monitoring

#### Reversible Causes


- Hypovolemia
- Hypoxia
- Hydrogen ion (acidosis)
- Hypoglycemia
- Hypo-/hyperkalemia
- Hypothermia
- Tension pneumothorax
- Tamponade, cardiac
- Toxins
- Thrombosis, pulmonary
- Thrombosis, coronary





## Pediatric Advanced Life Support

### Management of Shock After ROSC



### Estimation of Maintenance Fluid Requirements

- **Infants <10 kg:** 4 mL/kg per hour  
*Example:* For an 8-kg infant, estimated maintenance fluid rate  
= 4 mL/kg per hour × 8 kg  
= 32 mL per hour
- **Children 10-20 kg:** 4 mL/kg per hour for the first 10 kg + 2 mL/kg per hour for each kg above 10 kg  
*Example:* For a 15-kg child, estimated maintenance fluid rate  
= (4 mL/kg per hour × 10 kg)  
+ (2 mL/kg per hour × 5 kg)  
= 40 mL/hour + 10 mL/hour  
= 50 mL/hour
- **Children >20 kg:** 4 mL/kg per hour for the first 10 kg + 2 mL/kg per hour for kg 11-20 + 1 mL/kg per hour for each kg above 20 kg.  
*Example:* For a 28-kg child, estimated maintenance fluid rate  
= (4 mL/kg per hour × 10 kg)  
+ (2 mL/kg per hour × 10 kg)  
+ (1 mL/kg per hour × 8 kg)  
= 40 mL per hour + 20 mL per hour  
+ 8 mL per hour  
= 68 mL per hour

Following initial stabilization, adjust the rate and composition of intravenous fluids based on the patient's clinical condition and state of hydration. In general, provide a continuous infusion of a dextrose-containing solution for infants. Avoid hypotonic solutions in critically ill children; for most patients use isotonic fluid such as normal saline (0.9% NaCl) or lactated Ringer's solution with or without dextrose, based on the child's clinical status.



# Pediatric Color-Coded Length-Based Resuscitation Tape

## Pediatric Advanced Life Support



Equipment	GRAY* 3-5 kg	PINK Small Infant 6-7 kg	RED Infant 8-9 kg	PURPLE Toddler 10-11 kg	YELLOW Small Child 12-14 kg	WHITE Child 15-18 kg	BLUE Child 19-23 kg	ORANGE Large Child 24-29 kg	GREEN Adult 30-36 kg
Resuscitation bag		Infant/child	Infant/child	Child	Child	Child	Child	Child	Adult
Oxygen mask (NRB)		Pediatric	Pediatric	Pediatric	Pediatric	Pediatric	Pediatric	Pediatric	Pediatric/adult
Oral airway (mm)		50	50	60	60	60	70	80	80
Laryngoscope blade (size)		1 Straight	1 Straight	1 Straight	2 Straight	2 Straight	2 Straight or curved	2 Straight or curved	3 Straight or curved
ET tube (mm) <sup>†</sup>		3.5 Uncuffed 3.0 Cuffed	3.5 Uncuffed 3.0 Cuffed	4.0 Uncuffed 3.5 Cuffed	4.5 Uncuffed 4.0 Cuffed	5.0 Uncuffed 4.5 Cuffed	5.5 Uncuffed 5.0 Cuffed	6.0 Cuffed	6.5 Cuffed
ET tube insertion length (cm)	3 kg 9-9.5 4 kg 9.5-10 5 kg 10-10.5	10.5-11	10.5-11	11-12	13.5	14-15	16.5	17-18	18.5-19.5
Suction catheter (F)		8	8	10	10	10	10	10	10-12
BP cuff	Neonatal #5/infant	Infant/child	Infant/child	Child	Child	Child	Child	Child	Small adult
IV catheter (ga)		22-24	22-24	20-24	18-22	18-22	18-20	18-20	16-20
IO (ga)		18/15	18/15	15	15	15	15	15	15
NG tube (F)		5-8	5-8	8-10	10	10	12-14	14-18	16-18
Urinary catheter (F)	5	8	8	8-10	10	10-12	10-12	12	12
Chest tube (F)		10-12	10-12	16-20	20-24	20-24	24-32	28-32	32-38

Abbreviations: BP, blood pressure; ET, endotracheal; F, French; IO, intraosseous; IV, intravenous; NG, nasogastric; NRB, nonrebreathing.

\*For Gray column, use Pink or Red equipment sizes if no size is listed.

<sup>†</sup>Per 2010 AHA Guidelines, in the hospital cuffed or uncuffed tubes may be used (see below for sizing of cuffed tubes).

Adapted from Broselow™ Pediatric Emergency Tape. Distributed by Armstrong Medical Industries, Lincolnshire, IL. Copyright 2007 Vital Signs, Inc. All rights reserved.



